

The transient neutron star X-ray binary KS 1741–293 in outburst and quiescence

N. Degenaar¹† and R. Wijnands²

¹University of Michigan, Dept. of Astronomy, 500 Church St, Ann Arbor, MI 48109, USA
 email: degenaar@umich.edu

²Astronomical Institute “Anton Pannekoek”, University of Amsterdam
 Postbus 94249, 1090 GE Amsterdam, The Netherlands
 email: r.a.d.wijnands@uva.nl

Abstract. KS 1741–293 is a transient neutron star low-mass X-ray binary that is located at an angular distance of $\simeq 20'$ from the Galactic center. We map out the historic activity of the source since its discovery in 1989, characterize its most recent X-ray outbursts observed with *Swift* (2007, 2008, 2010, and 2011), and discuss its quiescent X-ray properties using archival *Chandra* data. KS 1741–293 is frequently active, exhibiting outbursts that typically reach a 2–10 keV luminosity of $L_X \simeq 10^{36} (D/6.2 \text{ kpc})^2 \text{ erg s}^{-1}$ and last for several weeks–months. However, *Swift* also captured a very short and weak accretion outburst that had a duration of $\lesssim 4$ days and did not reach above $L_X \simeq 5 \times 10^{34} (D/6.2 \text{ kpc})^2 \text{ erg s}^{-1}$. The source is detected in quiescence with *Chandra* at a 2–10 keV luminosity of $L_X \simeq 2.5 \times 10^{32} (D/6.2 \text{ kpc})^2 \text{ erg s}^{-1}$.

Keywords. accretion, accretion disks, stars: neutron, X-rays: binaries, X-rays: individual (KS 1741–293)

1. Introduction

KS 1741–293 is a transient neutron star low-mass X-ray binary (LMXB) that was discovered in 1989 August by the TTM onboard the KVANT module of the Mir space station (in 't Zand et al. 1991). The detection of type-I X-ray bursts revealed its binary nature and testified to the presence of a neutron star. The source is located at an angular distance of $\simeq 20'$ from the Galactic center, at an estimated distance of $D \simeq 6.2 \text{ kpc}$ (as inferred from type-I X-ray burst analysis; Chelovekov & Grebenev 2011). Throughout this work we assume a distance of $D \simeq 6.2 \text{ kpc}$ when quoting X-ray luminosities.

2. The outburst history of KS 1741–293

We compiled a list of all (non-) detections of KS 1741–293 reported in literature, and combined these with the analysis of archival data (Sections 3 and 4) to map out its historic activity. Fluxes and upper limits were converted to the 2–10 keV energy band using PIMMS, assuming a power law spectrum with $N_H = 21.5 \times 10^{22} \text{ cm}^{-2}$ and $\Gamma = 2.1$. The long-term X-ray light curve as observed between 1989–2012 is shown in Figure 1.

After its discovery in 1989, the field around KS 1741–293 was observed with *ROSAT* in 1992, but the source was not detected (Figure 1; Sidoli et al. 2001). Renewed activity was seen in 1998 March and September by *BeppoSAX* and *ASCA*, which possibly covered the same outburst (in 't Zand et al. 1998; Sidoli et al. 1999; Sakano et al. 2002). *INTEGRAL* detected it in outburst in 2003, 2004 and 2005 (de Cesare et al. 2007; Kuulkers et al. 2007; Chelovekov & Grebenev 2011). Non-detections in between these epochs suggest

† Hubble fellow




Figure 1. Long-term outburst history of KS 1741–293 (2–10 keV; see text for references).

that this were likely three separate outbursts (see Figure 1). The 2005 outburst was also observed by *Chandra* and *Suzaku* (Degenaar et al. 2008; Yuasa et al. 2008).

Between 2006 and 2008, the source region was covered by monitoring campaigns of the Galactic center with *Chandra* and *XMM-Newton* (Wijnands et al. 2006; Degenaar et al. 2012), and *Swift* (Degenaar & Wijnands 2009, 2010). The source was found (weakly) active on three different epochs in 2007 (Section 3; Degenaar et al. 2012), and a new outburst was observed in 2008 (Degenaar & Wijnands 2008; Degenaar et al. 2012). *INTEGRAL* and *Swift* found the source active again in 2010 (Chenevez et al. 2010) and in 2011 (Barthelmy et al. 2011; Linares et al. 2011; Chenevez et al. 2011).

Only a few type-I X-ray bursts have been reported for KS 1741–293. KVANT detected two (in ’t Zand et al. 1991), six have been seen with *INTEGRAL* (de Cesare et al. 2007; Chelovekov & Grebenev 2011), and recently one was picked up by *Swift* (Barthelmy et al. 2011; Linares et al. 2011). *RXTE* may have also detected one (Galloway et al. 2008).

3. Outbursts observed with *Swift* between 2007 and 2012

We obtained all *Swift*/XRT data covering KS 1741–293 from the public data archive. This encompasses 69 observations performed between 2007 May 5 and 2012 May 17. We extracted data products using the online XRT tools (Evans et al. 2009). Figure 2 displays the *Swift*/XRT light curve. It shows three major outbursts in 2008, 2010, and 2011, as well as a short and weak episode of activity in 2007. We extracted spectra for these four different outbursts and fitted these simultaneously in XSPEC to a simple absorbed power law model. The results of the spectral analysis are summarized in Table 1.

The region around KS 1741–293 was observed every few days by *Swift* between 2007 May 23 and August 9. The source is not detected during this epoch, except during three observations performed on June 13 and 14, when it displayed a 2–10 keV luminosity of $L_X \simeq 3 \times 10^{34} \text{ erg s}^{-1}$. Non-detections in the preceding and subsequent observations (June 11 and 15) suggest that this episode of low-level activity had a duration of $\lesssim 4$ days.

KS 1741–293 was detected at similarly low intensities at another two epochs in 2007. *XMM-Newton* found the source active at $L_X \simeq 8 \times 10^{34} \text{ erg s}^{-1}$ on 2007 September 6. Non-detections with *Swift* on August 9 and September 27 constrain the duration of this




Figure 2. Long-term *Swift*/XRT light curve (binned per observation) showing three main outbursts of KS 1741–293 (2008, 2010, and 2011) and one episode of low-level activity (2007).

outburst to be $\lesssim 49$ days. In an archival *Chandra* observation performed on 2007 April 25, KS 1741–293 is weakly detected at $L_X \simeq 1.3 \times 10^{33} \text{ erg s}^{-1}$ (Table 1). Non-detections with *Chandra* on 2007 February 19 ($L_X \lesssim 5 \times 10^{32} \text{ erg s}^{-1}$) and *Swift* between 2007 May 23 and June 10 ($L_X \lesssim 1 \times 10^{33} \text{ erg s}^{-1}$) suggest that the activity lasted for $\lesssim 3$ months. KS 1741–293 was thus found active a factor $\simeq 10$ –100 above its quiescent level several times in 2007, but was not detected above $L_X \simeq 10^{35} \text{ erg s}^{-1}$.

We obtained a series of *Swift*/XRT ToO observations of KS 1741–293 between 2008 May 18 and September 4 to monitor the new outburst that was first detected by *Chandra* on May 10 (Degenaar et al. 2008; Degenaar & Wijnands 2008). The source remained active for $\simeq 3$ months at an average 2–10 intensity of $L_X \simeq 8 \times 10^{35} \text{ erg s}^{-1}$ (Table 1) till August 21. It was no longer detected on September 4 with an upper limit of $L_X \lesssim 9 \times 10^{33} \text{ erg s}^{-1}$. In 2010 and 2011, KS 1741–293 was again observed in outburst with *Swift* at an intensity of $L_X \simeq (3 - 7) \times 10^{35} \text{ erg s}^{-1}$ (Table 1, Figure 2).

4. KS 1741–293 in quiescence

To investigate the quiescent properties, we used archival *Chandra* observations performed in 2001, 2006 and 2007 (Obs IDs 2267, 7038, and 8459; from the campaign of Munro et al. 2009). The source is not detected in the individual images, but clearly visible when the observations are combined ($\simeq 51 \text{ ks}$). We created a combined spectrum using the CIAO tools, and fitted this simultaneously with the *Swift* outburst data (Table 1).

KS 1741–293 is detected in quiescence at a 2–10 keV luminosity of $L_X \simeq 2.5 \times 10^{32} \text{ erg s}^{-1}$, which is typical for quiescent neutron star LMXBs. The very large extinction in the direction of the source ($N_H \simeq 2 \times 10^{23} \text{ cm}^{-2}$) completely obscures the thermal emission that is often detected for quiescent neutron star LMXBs (typically $kT_{\text{bb}} \simeq 0.1$ –0.3 keV). Therefore, we only detect the hard (non-thermal) power law tail.

5. Discussion

KS 1741–293 is a frequently active: between 1989 and 2012, the source exhibited at least 8 accretion outbursts that reached $L_X \simeq 10^{36} \text{ erg s}^{-1}$ (2–10 keV) and had a duration

Table 1. Spectral properties of KS 1741–293 in outburst and quiescence.

Instr.	Date	State	Γ	F_X (erg cm ⁻² s ⁻¹)	L_X (erg s ⁻¹)
<i>Chandra</i>	2001/2006/2007	quiescence	$1.3^{+5.4}_{-1.3}$	$(5.5 \pm 5.0) \times 10^{-14}$	$(2.5 \pm 2.3) \times 10^{32}$
<i>Chandra</i>	2007 Apr 25	low activity	$0.2^{+1.6}_{-0.2}$	$(2.8 \pm 1.3) \times 10^{-13}$	$(1.3 \pm 0.6) \times 10^{33}$
<i>Swift</i>	2007 Jun 13–14	low activity	$0.8^{+2.4}_{-0.8}$	$(6.2 \pm 1.4) \times 10^{-12}$	$(2.9 \pm 0.6) \times 10^{34}$
<i>Swift</i>	2008 May 18–Aug 21	outburst	2.1 ± 0.4	$(1.7 \pm 0.4) \times 10^{-10}$	$(7.8 \pm 1.9) \times 10^{35}$
<i>Swift</i>	2010 Mar 10	outburst	2.2 ± 0.6	$(1.5 \pm 0.5) \times 10^{-10}$	$(6.9 \pm 2.3) \times 10^{35}$
<i>Swift</i>	2011 Sep 1–30	outburst	2.7 ± 1.6	$(6.0 \pm 2.4) \times 10^{-11}$	$(2.8 \pm 1.1) \times 10^{35}$

Note. Quoted errors refer to 90% confidence levels. A simultaneous fit to the spectral data resulted in $N_H = (21.5 \pm 2.8) \times 10^{22}$ cm⁻² and $\chi^2_\nu = 1.1$ for 142 dof. F_X represents the unabsorbed 2–10 keV flux and L_X the corresponding luminosity assuming $D = 6.2$ kpc.

of several weeks–months. This suggests a recurrence time of $\simeq 2$ yr and a duty cycle of $\simeq 12.5\%$. In addition to these main outbursts, we found indications of low-level accretion activity a factor of $\simeq 10$ – 100 above the quiescent level of $L_X \simeq 2.5 \times 10^{32}$ erg s⁻¹. Such peculiar behavior has now been observed for a number of transient neutron star LMXBs, such as XMM J174457–2850.3, GRO J1744–28, GRS 1741–2853, and XTE J1701–462 (Degenaar & Wijnands 2009, 2010; Degenaar et al. 2012; Fridriksson et al. 2011).

Acknowledgements

ND is supported by NASA through Hubble Postdoctoral Fellowship grant number HST-HF-51287.01-A from the Space Telescope Science Institute, and RW by a European Research Council starting grant. This work made use of *Swift* data supplied by the UK Swift Science Data Centre at the University of Leicester, and the *Chandra* data archive.

References

- Barthelmy, S. D., Campana, S., D’Elia, V., et al. 2011, *GRB Coordinates Network*, 12319
- Chelovekov, I. V. & Grebenev, S. A. 2011, *Astronomy Letters*, 37, 597
- Chenevez, J., Brandt, S., Kuulkers, E., et al. 2011, *The Astronomer’s Telegram*, 3646
- Chenevez, J., Kuulkers, E., Alfonso-Garzón, J., et al. 2010, *The Astronomer’s Telegram*, 2465
- de Cesare, G., Bazzano, A., Martínez Núñez, S., et al. 2007, *MNRAS*, 380, 615
- Degenaar, N. & Wijnands, R. 2008, *The Astronomer’s Telegram*, 1541
- Degenaar, N. & Wijnands, R. 2009, *A&A*, 495, 547
- Degenaar, N. & Wijnands, R. 2010, *A&A*, 524, A69
- Degenaar, N., Wijnands, R., Cackett, E. M., et al. 2012, *A&A*, 545, A49
- Degenaar, N., Wijnands, R., & Munro, M. 2008, *The Astronomer’s Telegram*, 1531
- Evans, P. A., Beardmore, A. P., Page, K. L., et al. 2009, *MNRAS*, 397, 1177
- Fridriksson, J. K., Homan, J., Wijnands, R., et al. 2011, *ApJ*, 736, 162
- Galloway, D. K., Munro, M. P., Hartman, et al. 2008, *ApJS*, 179, 360
- in ’t Zand, J., Heise, J., Smith, M., et al. 1998, *IAU Circ.*, 6840, 2
- in ’t Zand, J. J. M., Heise, J., Brinkman, A. C., et al. 1991, *Adv. Space Res.*, 11, 187
- Kuulkers, E., Shaw, S. E., Paizis, A., et al. 2007, *A&A*, 466, 595
- Linares, M., Chakrabarty, D., Cummings, J. R., et al. 2011, *The Astronomer’s Telegram*, 3632
- Munro, M. P., Bauer, F. E., Baganoff, F. K., et al. 2009, *ApJS*, 181, 110
- Sakano, M., Koyama, K., Murakami, et al. 2002, *ApJS*, 138, 19
- Sidoli, L., Belloni, T., & Mereghetti, S. 2001, *A&A*, 368, 835
- Sidoli, L., Mereghetti, S., Israel, G. L., et al. 1999, *ApJ*, 525, 215
- Wijnands, R., in ’t Zand, J. J. M., Rupen, M., et al. 2006, *A&A*, 449, 1117
- Yuasa, T., Tamura, K., Nakazawa, K., et al. 2008, *PASJ*, 60, 207